Issue No. 56

January 2008

DISTANT EKOs

The Kuiper Belt Electronic Newsletter

Edited by: Joel Wm. Parker

ekonews@boulder.swri.edu

www.boulder.swri.edu/ekonews

CONTENTS

NEWS & ANNOUNCEMENTS

There were 7 new TNO discoveries announced since the previous issue of *Distant EKOs*: 2007 RM283, 2007 RT15, 2007 TC418, 2007 TD418, 2007 TZ417, 2007 VJ302, 2007 VK302 and 4 new Centaur/SDO discoveries: 2007 RG283, 2007 RH283, 2007 TA418, 2007 TB418 Reclassified objects: 2007 RW10 (NTrojan \rightarrow Centaur) Objects recently assigned numbers: 2000 GE147 = 1687002000 GP183 = 1687032001 FR185 = 1690712003 MW12 = 1745672004 PF115 = 175113Current number of TNOs: 1074 (including Pluto) Current number of Centaurs/SDOs: 213 Current number of Neptune Trojans: 5 Out of a total of 1292 objects: 555 have measurements from only one opposition 540 of those have had no measurements for more than a year 278 of those have arcs shorter than 10 days (for more details, see: http://www.boulder.swri.edu/ekonews/objects/recov_stats.gif)

PAPERS ACCEPTED TO JOURNALS

Origin of the Structure of the Kuiper Belt during a Dynamical Instability in the Orbits of Uranus and Neptune

H. Levison¹, A. Morbidelli², C. Van Laerhoven³, R. Gomes⁴, and K. Tsiganis⁵

¹ Southwest Research Institute, USA

- ² Observatoire de la Côte d'Azur, France
- ³ Department of Physics and Astronomy, University of British Colombia, Canada

⁴ Observatório Nacional/MCT, Brazil

⁵ Department of Physics, Aristotle University of Thessaloniki, Greece

We explore the origin and orbital evolution of the Kuiper belt in the framework of a recent model of the dynamical evolution of the giant planets, sometimes known as the *Nice* model. This model is characterized by a short, but violent, instability phase, during which the planets were on large eccentricity orbits. It successfully explains, for the first time, the current orbital architecture of the giant planets (Tsiganis et al. 2005, Nature 435, 459), the existence of the Trojans populations of Jupiter and Neptune (Morbidelli et al. 2005, Nature 435, 462), and the origin of the late heavy bombardment of the terrestrial planets (Gomes et al. 2005, Nature 435, 466). One characteristic of this model is that the proto-planetary disk must have been truncated at roughly 30 to 35 AU so that Neptune would stop migrating at its currently observed location. As a result, the Kuiper belt would have initially been empty.

In this paper we present a new dynamical mechanism which can deliver objects from the region interior to ~ 35 AU to the Kuiper belt without excessive inclination excitation. In particular, we show that during the phase when Neptune's eccentricity is large, the region interior to its 1:2 mean motion resonance becomes unstable and disk particles can diffuse into this area. In addition, we perform numerical simulations where the planets are forced to evolve using fictitious analytic forces, in a way consistent with the direct N-body simulations of the Nice model. Assuming that the last encounter with Uranus delivered Neptune onto a low-inclination orbit with a semi-major axis of ~ 27 AU and an eccentricity of ~ 0.3 , and that subsequently Neptune's eccentricity damped in ~ 1 My, our simulations reproduce the main observed properties of the Kuiper belt at an unprecedented level. In particular, our results explain, at least qualitatively: 1) the so-existence of resonant and non-resonant populations, 2) the eccentricity–inclination distribution of the Plutinos, 3) the peculiar semi-major axis – eccentricity distribution in the classical belt, 4) the outer edge at the 1:2 mean motion resonance with Neptune, 5) the bi-modal inclination distribution of the classical population, 6) the correlations between inclination and physical properties in the classical Kuiper belt, the existence of the so-called extended scattered disk. Nevertheless, we observe in the simulations a deficit of nearly-circular objects in the classical Kuiper belt.

To appear in: Icarus

For preprints, contact hal@boulder.swri.edu or on the web at http://arxiv.org/abs/0712.0553

An Outer Planet beyond Pluto and Origin of the Trans-Neptunian Belt Architecture

Patryk Sofia Lykawka¹ and Tadashi Mukai¹

¹ Kobe University, Department of Earth and Planetary Sciences, 1-1 rokkodai-cho, nada-ku, Kobe, 657-8501. Japan

Trans-Neptunian objects (TNOs) are remnants of a collisionally and dynamically evolved planetesimal disk in the outer solar system. This complex structure, known as the trans-Neptunian belt (or Edgeworth-Kuiper belt), can reveal important clues about disk properties, planet formation, and other evolutionary processes. In contrast to the predictions of accretion theory, TNOs exhibit surprisingly large eccentricities, e, and inclinations, i, which can be grouped into distinct dynamical classes. Several models have addressed the origin and orbital evolution of TNOs, but none have reproduced detailed observations, e.g., all dynamical classes and peculiar objects, or provided insightful predictions. Based on extensive simulations of planetesimal disks with the presence of the four giant planets and massive planetesimals, we propose that the orbital history of an outer planet with tenths of Earth's mass can explain the trans-Neptunian belt orbital structure. This massive body was likely scattered by one of the giant planets, which then stirred the primordial planetesimal disk to the levels observed at 40–50 AU and truncated it at about 48 AU before planet migration. The outer planet later acquired an inclined stable orbit (>100 AU; 20–40 deg) because of a resonant interaction with Neptune (an r:1 or r:2 resonance possibly coupled with the Kozai mechanism), guaranteeing the stability of the trans-Neptunian belt. Our model consistently reproduces the main features of each dynamical class with unprecedented detail; it also satisfies other constraints such as the current small total mass of the trans-Neptunian belt and Neptune's current orbit at 30.1 AU. We also provide observationally testable predictions.

To appear in: The Astronomical Journal

For preprints, contact patryk@dragon.kobe-u.ac.jp or on the web at http://harbor.scitec.kobe-u.ac.jp/~patryk/index-en.html

.....

High Precision Photometry of Extreme KBO $2003 EL_{61}$

P. Lacerda¹, David Jewitt¹, and Nuno Peixinho^{1,2}

¹ Institute for Astronomy, University of Hawaii, USA ² Grupo de Astrofísica, Universidade de Coimbra, Portugal

We present high precision, time-resolved visible and near infrared photometry of the large (diameter ~ 2500 km) Kuiper belt object (136108) 2003 EL₆₁. The new data confirm rapid rotation at period $P = 3.9155\pm0.0001$ hr with a peak-to-peak photometric range $\Delta m_R = 0.29\pm0.02$ mag. and further show subtle but reproducible color variations with rotation. Rotational deformation of 2003 EL₆₁ alone would give rise to a symmetric lightcurve free of color variations. The observed photometric deviations from the best-fit equilibrium model show the existence of a large surface region with an albedo and color different from the mean surface of 2003 EL₆₁. We explore constraints on the nature of this anomalous region set by the existing data.

To appear in: The Astrophysical Journal For preprints, contact pedro@ifa.hawaii.edu or on the web at http://www.ifa.hawaii.edu/~pedro/papers.html

Surface Composition and Temperature of the TNO Orcus

M.A. Barucci¹, F. Merlin¹, A. Guilbert¹, C. de Bergh¹, A. Alvarez-Candal¹, O. Hainaut²,

A. Doressoundiram 1 , C. Dumas 2 , T. Owen 3 , and A. Coradini 4

 1 LESIA, Observatoire de Paris, F-92195 Meudon Principal Cedex, France

 2 ESO, Alonso de Cordova 3107, Vitacura, Casilla 19001, Santiago 19, Chile

 3 Institute or Astronomy 2680 Woodlawn Drive, Honolulu, HI
96822, USA

 4 INAF-IFSI, Via del Fosso del Cavaliere, 00133 Roma, Italy

The aim of this paper is to investigate the surface composition of the Transneptunian Object (TNO) Orcus. High quality observations have been carried out with the new instrument SINFONI at the Very Large Telescope (VLT) of ESO. Crystalline water ice, and possibly ammonia ice, have been found from spectroscopic observations of the TNO Orcus between 1.4 and 2.4 μ m. The existence of such ices on the surface of Orcus may indicate a renewal mechanism on the surface and geological activity. The presence of ammonia on the surface of Orcus, if confirmed, could have important implications for the composition of the primitive solar nebula and the formation of the TNO population.

To appear in: Astronomy & Astrophysics

For preprints, contact antonella.barucci@obspm.fr

Evidence of N_2 -Ice On the Surface of the Icy Dwarf Planet 136472 (2005 FY9)

S.C. Tegler¹, W.M. Grundy², F. Vilas³,

W. Romanishin⁴, D. Cornelison¹, and G.J. Consolmagno⁵

 1 Department of Physics and Astronomy, Northern Arizona University, Flagstaff, AZ, 86011, USA

 2 Lowell Observatory, 1400 W. Mars Hill Rd, Flagstaff, AZ, 86001, USA

³ MMT Observatory, PO Box 210065, University of Arizona, Tucson, AZ, 85721, USA

⁴ Department of Physics and Astronomy, University of Oklahoma, Norman, OK, 73019, USA

.....

⁵ Vatican Observatory, Specola, Vaticana, V-00120, Vatican City State

We present high signal precision optical reflectance spectra of 2005 FY9 taken with the Red Channel Spectrograph and the 6.5-m MMT telescope on 2006 March 4 UT (5000–9500 Å; 6.33 Å pixel⁻¹) and 2007 February 12 UT (6600–8500 Å; 1.93 Å pixel⁻¹). From cross correlation experiments between the 2006 March 4 spectrum and a pure CH₄-ice Hapke model, we find the CH₄-ice bands in the MMT spectrum are blueshifted by 3 ± 4 Å relative to bands in the pure CH₄-ice Hapke spectrum. The higher resolution MMT spectrum of 2007 February 12 UT enabled us to measure shifts of individual CH₄-ice bands. We find the 7296 Å, 7862 Å, and 7993 Å CH₄-ice bands are blueshifted by 4 ± 2 Å, 4 ± 4 Å, and 6 ± 5 Å. From four measurements we report here and one of our previously published measurements, we find the CH₄-ice bands in spectra of 2005 FY9. Laboratory experiments show that CH₄-ice bands in spectra of CH₄ mixed with other ices are blueshifted relative to bands in spectra of pure CH₄-ice. A likely candidate for the other component is N₂-ice because its weak 2.15 μ m band and blueshifted CH₄ bands are seen in spectra of Triton and Pluto. Assuming the shift is due to the presence of N₂, spectra taken on two consecutive nights show no difference in CH₄/N₂. In addition, we find no measurable difference in CH₄/N₂ at different depths into the surface of 2005 FY9.

To appear in: Icarus

For preprints, contact Stephen.Tegler@nau.edu or on the web at http://arxiv.org/abs/0801.3115

A Search for sub-km KBOs with the Method of Serendipitous Stellar Occultations

S.J. Bickerton¹, JJ. Kavelaars², and D.L. Welch³

¹ McMaster University, Dept. of Physics and Astronomy, 1280 Main St. West, Hamilton, ON L8S 4M1, Canada

² Herzberg Institute of Astrophysics, 5071 West Saanich Rd., Victoria, BC V9E 2E7, Canada

³ McMaster University, Dept. of Physics and Astronomy, 1280 Main St. West, Hamilton, ON L8S 4M1, Canada

The results of a search for sub-km Kuiper Belt Objects (KBOs) with the method of serendipitous stellar occultations are reported. Photometric time series were obtained on the 1.8m telescope at the Dominion Astrophysical Observatory (DAO) in Victoria, British Columbia, and were analyzed for the presence of occultation events. Observations were performed at 40 Hz and included a total of 5.0 star-hours for target stars in the ecliptic open cluster M35 ($\beta = 0.9^{\circ}$), and 2.1 star-hours for control stars in the off-ecliptic open cluster M34 ($\beta = 25.7^{\circ}$). To evaluate the recovery fraction of the analysis method, and thereby determine the limiting detectable size, artificial occultation events were added to simulated time series (1/f scintillation-like power-spectra), and to the real data. No viable candidate occultation events were detected. This limits the cumulative surface density of KBOs to $3.5 \times 10^{10} \text{ deg}^{-2}$ (95% confidence) for KBOs brighter than m_R=35.3 (larger than ~860 m in diameter, assuming a geometric albedo of 0.04 and a distance of 40 AU). An evaluation of TNO occultations reported in the literature suggests that they are unlikely to be genuine, and an overall 95%-confidence upper limit on the surface density of 2.8 × 10⁹ deg⁻² is obtained for KBOs brighter than m_R=35 (larger than ~1 km in diameter, assuming a geometric albedo of 0.04 and a distance of 40 AU) when all existing surveys are combined.

To appear in: The Astronomical Journal

For preprints, contact bick@astro.princeton.edu or on the web at http://arXiv.org/abs/0801.2969

Helios and Prometheus: A Solar/Nuclear Outer-Solar System Mission

.....

Gregory L. Matloff¹, Les Johnson², and Claudio Maccone³

¹ New York City College of Technology, CUNY, USA

² Marshall Space flight Center, NASA, USA

³ Politechnico Torino, Turin, Italy

A 2–3 decade mission is proposed with a solar-sail spacecraft approaching the Sun within 0.2 AU. After sail unfurlment at the perihelion of an initially elliptical solar orbit and the completion of acceleration, the spacecraft splits into two components. One part is a scientific payload bound for the heliopause; the second is designed to rendezvous with a Kuiper Belt Object and decelerates across the solar system using Radioisotope-Electric Propulsion.

Published in: Journal of the British Interplanetary Society, 60, 439 (2007 Dec) For preprints, contact GMatloff@citytech.cuny.edu

PAPERS RECENTLY SUBMITTED TO JOURNALS

Masses of Nix and Hydra

David J. Tholen¹, Marc W. Buie², William M. Grundy², and Garrett T. Elliott³

¹ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

 2 Lowell Observatory, 1400 W. Mars Hill Road, Flagstaff, AZ 86001, USA

 3 The Ohio State University, Columbus, OH 43210, USA

Submitted to: The Astronomical Journal

Preprint available on the web at http://arxiv.org/abs/0712.1261

OTHER PAPERS OF INTEREST

Amorphization of Crystalline Water Ice Weijun Zheng^{1,2,3}, David Jewitt¹, and Ralf I. Kaiser²

¹ Institute for Astronomy, University of Hawaii, Honolulu, HI 96822, USA

² Department of Chemistry, University of Hawaii, Honolulu, HI 96822, USA

³ State Key Laboratory of Molecular Reaction Dynamics, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100080, P. R. China

Preprint available on the web at http://arxiv.org/abs/0801.2805

CONFERENCE CONTRIBUTIONS

Detection of Small Kuiper Belt Objects by Stellar Occultations ${{\scriptstyle R.Stevenson}^1}$

¹ Institute for Astronomy, 2680 Woodlawn Drive, University of Hawaii, Honolulu, HI 96822, USA

To appear in: Proc. of the 14th Young Scientists Conference on Astronomy and Space Physics Preprint available on the web at http://arxiv.org/abs/0712.1550

BOOKS

The Solar System Beyond Neptune

Edited by M. Antonietta Barucci Hermann Boehnhardt Dale P. Cruikshank Alessandro Morbidelli

The University of Arizona Press in collaboration with the Lunar and Planetary Institute, 2008

CONTENTS

PART I: INTRODUCTION
The Solar System Beyond Neptune: Overview and Perspectives
M.A. Barucci, H. Boehnhardt, D.P. Cruikshank, and A. Morbidelli
The Early Development of Ideas Concerning the Transneptunian Region
J.K. Davies, J. McFarland, M.E. Bailey, B.G. Marsden, and WH. Ip
Transneptunian Orbit Computation
J. Virtanen, G. Tancredi, G.M. Bernstein, T. Spahr, and K. Muinonen
PART II: TRANSNEPTUNIAN OBJECT POPULATIONS
Nomenclature in the Outer Solar System
B. Gladman, B.G. Marsden, and C. VanLaerhoven
The Orbital and Spatial Distribution of the Kuiper Belt
JJ Kavelaars, L. Jones, B. Gladman, J.W. Parker, and JM. Petit
Size Distribution of Multikilometer Transneptunian Objects
J-M. Petit, JJ Kavelaars, B. Gladman, and T. Laredo
PART III: BULK PROPERTIES
Color Properties and Trends of the Transneptunian Objects
A. Doressoundiram, H. Boehnhardt, S.C. Tegler, and C. Trujillo
Colors of Centaurs
S.C. Tegler, J.M. Bauer, W. Romanishin, and N. Peixinho
Surface Properties of Kuiper Belt Objects and Centaurs from Photometry and Polarimetry
I.N. Belskaya, A-C. Levasseur-Regourd, Y.G. Shkuratov, and K. Muinonen
Photometric Lightcurves of Transneptunian Objects and Centaurs: Rotations, Shapes, and Densities
S.S. Sheppard, P. Lacerda, and J.L. Ortiz
Composition and Surface Properties of Transneptunian Objects and Centaurs
M.A. Barucci, M.E. Brown, J.P. Emery, and F. Merlin
Physical Properties of Kuiper Belt and Centaur Objects: Constraints from the Spitzer Space Telescope
J. Stansberry, W. Grundy, M. Brown, D. Cruikshank, J. Spencer, D. Trilling, and JL. Margot 161
Transneptunian Object Taxonomy
M. Fulchignoni, I. Belskaya, M.A. Barucci, M.C. De Santis, and A. Doressoundiram

5
~
3
2
ა
0
9
5
0
3
5
5
5
-
Э
~
3
7
1
1
5
3
5
n
0
3
7
'

PART X: PERSPECTIVES

The Kuiper Belt Explored by Serendipitous Stellar Occultations	
F. Roques, G. Georgevits, and A. Doressoundiram	545
New Horizons: NASA's Pluto-Kuiper Belt Mission	
H.A. Weaver and S.A. Stern	557
Future Surveys of the Kuiper Belt	
C.A. Trujillo	573
0.11. 1. 4)	0.0

THESES

A Search for Kilometer-Sized Kuiper Belt Objects with the Method of Serendipitous Stellar Occultations

S.J. Bickerton¹

¹ McMaster University, Dept. of Physics and Astronomy, 1280 Main St. West, Hamilton, ON L8S 4M1, Canada

The results of a search for kilometer-sized Kuiper Belt Objects (KBOs) with the method of serendipitous stellar occultations are reported. Photometric time-series were obtained at the Dominion Astrophysical Observatory (DAO) in Victoria, British Columbia, and were analyzed for the presence of occultation events. Observations were performed at 40 Hz and included a total of 5.0 star-hours for target stars in the ecliptic open cluster M35, and 2.1 star-hours for control stars in the off-ecliptic open cluster M34. To evaluate the recovery fraction of the analysis method, and thereby determine the limiting detectable size, artificial occultation events were added to the data. No viable candidate occultation events were detected. This limits the cumulative surface density of KBOs to $3.5 \times 10^{10} \text{ deg}^{-2}$ (95% confidence) for KBOs brighter than m_R=35.3 (larger than ~860 m in diameter, assuming a geometric albedo of 0.04 and a distance of 40 AU). To evaluate false-positive rates, a method of simulating scintillation effects was developed.

Dissertation directed by D.L. Welch and J-J. Kavelaars Ph.D. awarded May, 2007 from McMaster University Available in postscript or PDF, contact bick@astro.princeton.edu The *Distant EKOs* Newsletter is dedicated to provide researchers with easy and rapid access to current work regarding the Kuiper belt (observational and theoretical studies), directly related objects (e.g., Pluto, Centaurs), and other areas of study when explicitly applied to the Kuiper belt.

We accept submissions for the following sections:

- \star Abstracts of accepted papers
- * Titles of submitted (but not yet accepted) papers and conference articles
- \star Thesis abstracts
- \star Short articles, announcements, or editorials
- * Status reports of on-going programs
- \star Requests for collaboration or observing coordination
- \star Table of contents/outlines of books
- \star Announcements for conferences
- \star Job advertisements
- \star General news items deemed of interest to the Kuiper belt community

A IAT_EX template for submissions is appended to each issue of the newsletter, and is sent out regularly to the e-mail distribution list. Please use that template, and send your submission to:

ekonews@boulder.swri.edu

The *Distant EKOs* Newsletter is available on the World Wide Web at:

http://www.boulder.swri.edu/ekonews

Recent and back issues of the newsletter are archived there in various formats. The web pages also contain other related information and links.

Distant EKOs is not a refereed publication, but is a tool for furthering communication among people interested in Kuiper belt research. Publication or listing of an article in the newsletter or the web page does not constitute an endorsement of the article's results or imply validity of its contents. When referencing an article, please reference the original source; *Distant EKOs* is not a substitute for peer-reviewed journals.

Moving ... ??

If you move or your e-mail address changes, please send the editor your new address. If the newsletter bounces back from an address for three consecutive issues, the address will be deleted from the mailing list. All address changes, submissions, and other correspondence should be sent to:

ekonews@boulder.swri.edu